

A REMARKABLE NEW SPECIES OF CERATINELLA EMERTON , 1882
(ARANEAE, LINYPHIIDAE, ERIGONINAE) FROM THE RUSSIAN FAR
EAST

Yuri M. Marusik^{1,2} , Valery A. Gnelitsa^{2,3}

1. Institute for Biological Problems of the North , Portovaya Str. 18 , Magadan , 685000 Russia ; E-mail : yumar@mail.ru

2. Zoological Museum , University of Turku , FI-20014 Turku , Finland

3. Sumy State Makarenko Teacher's Training University , 87 Romenskaya Str. , Sumy 40000 , Ukraine ; E-mail : gnelitsa@mail.ru

Abstract The new species *Ceratinella kurenschikovi* sp. nov. is described on the basis of the holotype male from Khabarovsk Province. It has a modified carapace , a character unknown in other *Ceratinella* species. The male palp of the new species is most similar to that of *C. brevis*.

Key words Spider , cephalic outgrowth , scutum , Bolshekhokhtsyrski Reserve.

Introduction

Ceratinella is a relatively large , but poorly studied genus of erigonine spiders. It comprises of 28 recognized species and has a Holarctic distribution (Platnick , 2009) . Only one species , *C. sydneyensis* Wunderlich , 1976 is known outside of the Holarctic region. This species is known only from the female and is either misplaced or has been introduced to Australia. Of the 28 species 11 are known only from females , and two other species are known only from males. Of the 11 species described from females , 7 are from the Nearctic and all of them were described by R. Chamberlin. The copulatory organs in this genus are rather homogeneous and the epigyne is only slightly variable. Only two Nearctic species : *C. brunnea* Emerton , 1882 and *C. buna* Chamberlin , 1949 have modified palps and epigynes. Males of these species have long retrolateral tibial apophyses and females lack an apical hood (socket) of epigyne. Some species are definitely misplaced (*C. subulata* Bösenberg & Strand , 1906 and *C. fumifera* Saito , 1939) . Eight species of *Ceratinella* are known from Russia (Mikhailov , 1997) . Only one of them , *C. sibirica* Strand , 1903 , described from Krasnoyarsk , is known only from the female.

While studying material from Khabarovsk Province we found one male with a remarkably modified cephalic region , but with the palp and abdomen typical for *Ceratinella*. The aim of this paper is to describe this new species. The holotype male will be deposited in the Zoological Museum of the Moscow State University (ZMMU) .

The holotype was photographed using an Olympus Camedia C-5050 camera attached to an Olympus SZX12 stereomicroscope. The images were montaged using "Combine ZM " image stacking software. Photographs were taken in dishes with paraffin on the bottom. Holes were made in the bottom to keep specimens in the correct position. All measurements are given in millimetres.

Species description

Ceratinella kurenschikovi sp. nov. (Figs. 1-10)

Material. Holotype (ZMMU) , Russia , Khabarovsk Prov. , environs of Khabarovsk , Bolshekhokhtsyrski Reserve , Klyuch Sosninsky Kordon (48 14 N , 134 47 E) , broad leafed forest , ca 400 m , fallen tree , 8.06.1994 (D. K. Kurenschikov) .

Etymology. The specific name is a patronym in honour of our friend and colleague Dmitri K. Kurenschikov who collected the holotype.

Diagnosis. This species can be easily distinguished from other *Ceratinella* and even all other Erigoninae by the shape of the cephalic outgrowth (cf. Figs. 1-3 , 8-10) . In terms of male palp structure , the new species is very similar to *C. brevis* (Wider , 1834) (the type species of *Ceratinella*) , but can be easily differentiated by the shape of the carapace and the large abdominal scutum covering the entire dorsal surface.

Description. Total length 1.54. Carapace : 0.7 long , 0.66 wide , yellow-red to brown. Cephalic region modified , with T-shaped outgrowth bearing PME (Figs. 1-3 , 8-10) . PME separated by approximately 4 times their diameters. Sternum 0.43 long , 0.42 wide , orange. Labium dark brown. Tibial spines absent or indistinct.

Table 1. Leg joints length.

Femur	Patella	Tibia	Metatarsus	Tarsus	Mt Tr
0.53	0.19	0.43	0.36	0.32	0.47
0.53	0.18	0.43	0.37	0.30	0.50
0.43	0.18	0.32	0.31	0.25	0.46
0.59	0.18	0.50	0.42	0.29	0.49

Entire abdomen covered by light brown scutum with 4 sigillae , the posterior pair being more distinct than the anterior pair (Figs. 2-3) . Palp as in Figs. 4-7. Femur and patella unmodified ; patella with small claw-like



Figs. 1-3. General appearance of *Ceratinella kurenschikovi* sp. nov. 1. Frontal. 2. Dorsal. 3. Cadual. Scale bars = 0.5 mm.

retrolateral apophysis and rectangular dorsal apophysis , cymbium with small prolateral lobe (Fig. 5) , paracymbium massive , embolus proper thin , tail-piece of radix forming a sharp point.

Comment. Although no other *Ceratinella* species has a modified carapace and scutum covering the entire dorsal side of the abdomen , we placed the new species in this genus because of the conformation of the male palp , which is almost identical to *C. brevis* , the type species of the genus. The presence or absence of a cephalic outgrowth in Erigoninae cannot serve as a generic character , because in several genera there are some species in which males have a modified prosoma and other species that do not , i. e. *Ceraticelus* , *Oedothorax* , *Perlongipalpus* , *Procerocymbium* , *Walckenaeria* , etc.


According to Millidge (1977) *Ceratinella* belongs to the *Peleopsis* group of genera. The most closely related

genera are *Ceratinopsis* Emerton , 1882 , *Styloctetor* Simon , 1884 and *Ceraticelus* Simon , 1884. The first two genera have males with an unmodified carapace , and only one *Styloctetor* species has an abdominal scutum.

Most of the *Ceraticelus* species with a modified carapace in males , have also developed dorsal and ventral abdominal scuta. At the same time *Ceraticelus* have a long tibial apophysis and females lack an anterior hood (socket).

Distribution. Type locality only. Because Bolshekhetsyrski Reserve lies on the border with China its occurrence there is very likely.

Acknowledgements This work was supported in part by the RFFI grant # 09-04-01365- . The English of the final draft was kindly checked by David Penney , Manchester.



Figs. 4-10. Left male palp and prosoma of *Ceratinella kurenschikovi* sp. nov. 4. Palp, retrolateral. 5. Palp, ventral. 6. Palp, prolateral. 7. Tibial apophysis, dorsal. 8. Prosoma, dorsal. 9. Prosoma, lateral. 10. Prosoma, frontal. Scale bars = 0.1 mm.

REFERENCES

- Platnick, N. I. 2009. The World Spider Catalog, Version 9.5. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html> (Accessed 6 April 2009).
- Mikhailov, K. G. 1997. Catalogue of the spiders of the territories of the former Soviet Union (Arachnida, Aranei). Zoological Museum of the Moscow State University, Moscow. 416 pp.
- Millidge, A. F. 1977. The conformation of the male palpal organs of linyphiid spiders, and its application to the taxonomic and phylogenetic analysis of the family (Araneae: Linyphiidae). Bull. Br. Arachnol. Soc., 4: 1-60.